

Maria João Bastos Lobato de Sousa

The impact of Computed Tomography Enterography and
Magnetic Resonance Enterography on the outcome of patients
with Crohn's Disease / O impacto da Enterografia por
Tomografia Computorizada e por Ressonância Magnética no
outcome dos doentes com Doença de Crohn

março, 2017

Maria João Bastos Lobato de Sousa

The impact of Computed Tomography Enterography and Magnetic
Resonance Enterography on the outcome of patients with Crohn's Disease
/ O impacto da Enterografia por Tomografia Computorizada e por
Ressonância Magnética no *outcome* dos doentes com Doença de Crohn

Mestrado Integrado em Medicina

Área: Gastroenterologia e Radiologia

Tipologia: Dissertação

Trabalho efetuado sob a Orientação de:

Professor Doutor Fernando Magro

Trabalho organizado de acordo com as normas da revista:

Inflammatory Bowel Diseases

março, 2017

FMUP

Eu, Maria João Bastos Lobato de Sousa, abaixo assinado, nº mecanográfico 201108530, estudante do 6º ano do Ciclo de Estudos Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 19 / 03 / 2017

Assinatura conforme cartão de identificação:

Maria João Bastos Lobato de Sousa

Projecto de Opção do 6º ano – DECLARAÇÃO DE REPRODUÇÃO

NOME

Maria João Bastos Lobato de Sousa

NÚMERO DE ESTUDANTE

201108530

E-MAIL

mariajoao.bastos.8@gmail.com

DESIGNAÇÃO DA ÁREA DO PROJECTO

Radiologia e Gastroenterologia

TÍTULO DISSERTAÇÃO/MONOGRAFIA (riscar o que não interessa)

The impact of Computed Tomography Enterography and Magnetic Resonance Enterography on the outcome of patients with Crohn's Disease

ORIENTADOR

Professor Doutor Fernando Magro

COORIENTADOR (se aplicável)

ASSINALE APENAS UMA DAS OPÇÕES:

| | |
|---|-------------------------------------|
| É AUTORIZADA A REPRODUÇÃO INTEGRAL DESTES TRABALHOS APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE. | <input type="checkbox"/> |
| É AUTORIZADA A REPRODUÇÃO PARCIAL DESTES TRABALHOS (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE. | <input checked="" type="checkbox"/> |
| DE ACORDO COM A LEGISLAÇÃO EM VIGOR, (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) NÃO É PERMITIDA A REPRODUÇÃO DE QUALQUER PARTE DESTES TRABALHOS. | <input type="checkbox"/> |

Faculdade de Medicina da Universidade do Porto, 19/03/2017

Assinatura conforme cartão de identificação: Maria Joã Bastos Lobato de Sousa

The impact of Computed Tomography Enterography and Magnetic Resonance Enterography on the outcome of patients with Crohn's Disease

Author: Maria João Bastos Lobato de Sousa⁽¹⁾ and Fernando José Magro Dias⁽²⁾

(1) *Medical Student*. Faculty of Medicine of Porto. Porto, Portugal.

(2) *Doctor of Medicine. Doctor of Philosophy*. Gastroenterology. Centro Hospitalar de São João, Faculty of Medicine of Porto. Porto, Portugal. Department of Pharmacology and Therapeutics, Faculty of Medicine of Porto. Porto, Portugal.

Address correspondence to: F. Magro, Department of Pharmacology and Therapeutics, Faculty of Medicine of Porto, Rua Plácido Costa, 4200-319 Porto, Portugal (e-mail: fm@med.up.pt).

The authors have no conflict of interest to disclose.

Abstract

Background: In the last few decades, Computed Tomography Enterography (CTE) and Magnetic Resonance Enterography (MRE) have become crucial weapons for the evaluation of patients with Crohn's Disease (CD). Despite the many studies on the role of cross sectional imaging in the diagnosis and assessment of CD activity, there are few related to their impact on the patients' outcome.

Methods: The study group for this retrospective study was composed of 349 patients previously diagnosed with CD, who underwent CTE/MRE from January 2010 to December 2015. Epidemiological and clinical data were collected from patients' clinical reports as well as CTE and MRE findings and their influence on management plan. Chi-square test and Fisher's exact test were used to evaluate the statistical significance.

Results: 349 cases were evaluated, of which 85% underwent CTE and 15% MRE. Overall, 60 patients had no CTE/MRE evidence of active disease and 289 had abnormal exams, of which the most frequent findings included bowel wall thickening and mural hyperenhancement. The CTE and MRE studies influenced a change in management in 40% of patients, leading more frequently to the introduction of immunomodulators (10%), anti-TNF α agents (8%) and corticosteroids (6%). Mural stratification was the only sign shown to be strongly related to change in management.

Conclusions: CTE and MRE played a decisive role in the outcome, leading to a therapeutic change in 40% of patients, and mural stratification, a sign of gravity, was strongly related with the outcome, leading to change in management in 46% of patients. Our findings further support the use of this imaging techniques in CD management algorithms.

Key Words: "Crohn's Disease", "Magnetic Resonance", "Computed Tomographic", "Enterography" and "Management Plan".

Introduction

Crohn's Disease (CD) is a chronic inflammatory bowel disease with a peak incidence between the second and fourth decades of life, characterized by an unpredictable course, marked by relapses and remissions.¹ It can affect any part of the gastrointestinal tract, from the mouth to the anus, with about 80% of cases involving the small bowel, particularly the terminal ileum.² The inflammation manifests itself by ulceration, which can range from superficial lesions to deep ulcers, with areas of inflammation interspersed with regions of normal mucosa (skip lesions). The transmural nature of CD always leads to stricturing or penetrating disease with intestinal obstruction and formation of fistulas and abscesses.³ The annual incidence of Crohn's Disease is estimated at 12.7 per 100,000 people in Europe and 20.2 per 100,000 people in North America. Its increasing incidence in both developed and developing countries shows the emergency of CD as a global disease.⁴

Several clinical indices have been developed in order to assess and quantify the severity of the disease. The most commonly used are the CD Activity Index and the Harvey Bradshaw Index, which classify the disease from asymptomatic remission to severe disease.⁵ With the recent introduction of effective medication, such as anti-TNF agents, "mucosal healing" is considered as the target of therapy for CD. This requires full disease assessment based not only on clinical and endoscopic data, but also on the findings of imaging techniques.⁶

Ileocolonoscopy is the gold standard for direct visualization of the colon's mucosa and for obtaining material for histological study through biopsies and it remains the ideal method for objective evaluation of disease activity in CD.^{7,8} However, in addition to not being able to evaluate the trans-mural extent of the disease, another major limitation of endoscopy is the inability to classify the disease severity at the level of the small intestine or to evaluate extra-luminal complications.⁵ Thus, cross-sectional imaging is becoming indispensable in the evaluation and classification of the severity of CD, since it allows the imaging of the entire gastrointestinal tract, observation of the bowel wall and evaluation of extra-intestinal complications, providing additional information and enabling clinicians to make more informed decisions regarding to patients' management plans.⁹ In the last consensus of the European Crohn's and Colitis Organisation, Computed Tomography Enterography (CTE) and Magnetic Resonance

Enterography (MRE) are the imaging techniques recommended to determine the extent of CD in the small intestine.¹⁰

CTE is widely used in the diagnosis of CD and evaluation of disease activity. Nevertheless, this imaging technique resorts to ionizing radiation with an adverse cumulative effect for patients, which is of extreme importance in CD since it affects mainly young adults and has a course marked by frequent relapses. Therefore, a free-radiation alternative is preferred, as is the case of MRE.⁷ There have been several studies comparing the effectiveness of MRE and CTE for detecting active inflammatory disease. A recent meta-analysis including 290 patients from 6 different studies diagnosed with CD showed that CTE and MRE have comparable sensitivity and specificity for demonstrating disease activity and detecting CD's complications.¹¹

An accurate assessment of disease characteristics is essential for proper management of CD, with prognostic implications.¹² Treat-to-target strategies are viewed as essential to the goal of altering the natural history of CD.¹³ With the increased use of cross sectional imaging in diagnosis and monitoring of CD, MRE-based severity scores have been developed and discussed. These severity scores can provide a better description of inflammatory lesions and severity assessment and may be useful in clinical practice to guide treatment decisions and in the evaluation of therapeutic efficacy.¹³ Rimola et al¹⁴ proposed and validated the most widely used severity score - "Magnetic Resonance Index of Activity" (MaRIA). This score takes into account bowel wall thickness, mucosal hyperenhancement and presence of edema and ulcers.⁵ MaRIA score was shown to be highly correlated with the CD Endoscopic Index of Severity (CDEIS), the reference gold standard for endoscopic evidence of disease.¹³

Despite the many studies on the role of CTE and MRE in the diagnosis and detection of CD activity, there are few related to their impact on the outcome of these patients.³ The objectives of the present study were to evaluate how CTE and MRE influence the clinician's therapeutic decision in patients with CD and to assess the impact of specific radiological signs in this decision.

Materials and Methods

This paper is based on a retrospective study of 349 patients of *Centro Hospitalar de São João* (CHSJ), with an established diagnosis of CD and who underwent CTE or MRE from January 2010 to December 2015. Patients with “suspected CD” as an indication were not included in this analysis. Patients with a diagnosis of Indeterminate Colitis or Ulcerative Colitis were excluded from our study. The cases in which more than 4 months had elapsed between the CTE or MRE evaluation and the change in management plans were also excluded.

The list of patients who underwent CTE or MRE between January 2010 and December 2015 was provided by the Radiology Department of CHSJ's. The clinical variables analyzed in our study were collected from the database of the *Grupo de Estudos de Doença Inflamatória Intestinal* (GEDII).

Of the 923 patients with inflammatory bowel disease who underwent CTE and ERM between 2010 and 2015 at our institution, 454 were excluded due to insufficient data for the analyses; 28 were excluded due to their diagnosis of Ulcerative Colitis or Undetermined Colitis; 19 were excluded because there was more than 4 months between the CTE or MRE assessment and the change in management, and 73 were excluded because these exams were performed with the purpose of aiding the diagnosis of CD.

Data Collection

Clinical and Demographic Data

Clinical data related to the Montreal Classification, which includes age at the diagnosis (A1 - ≤ 16 years old; A2 - 17-39 years old; and A3 - ≥ 40 years old); disease behavior (B1 – non stricturing non penetrating; B2 – stricturing; e B3 – penetrating; and p - perianal disease) and disease location (L1 - ileal; L2 - colonic; L3 - ileocolonic; and L4 – upper gastrointestinal tract), were collected. Clinically, this classification of CD has benefits in terms of patient counseling, assessment of disease prognosis, and providing support for the clinicians' decision of the most appropriate therapeutic plan for each disease subtype.^{15,16}

Other data obtained included clinical data related to a family history of CD, clinical occurrences during the course of the disease ("Emergency Service", "Hospitalizations" and "Surgeries"), reasons and clinical circumstances of CTE or MRE evaluation ("Acute disease" or "Follow-up"; "Hospitalization" or "Ambulatory") and previous upper and/or low digestive endoscopy.

The baseline characteristics of the patients are represented in table 1.

Prior Endoscopic Study

In an attempt to avoid possible detection biases, previous endoscopic studies were evaluated, estimating their possible influence on patient outcome. We took into account the studies that were included in the consultation report, in which CTE or MRE were evaluated.

CTE and MRE Reports

Any signs pertaining to disease activity were collected based on the original CTE and MRE reports (multiple radiologists) by one of the authors, and And the doubts were discussed with the other authors. The CTE and MRE findings recorded included: "Bowel wall thickening", "Mucosal enhancement", "Mural stratification", "Comb sign", "Ulcer", "Fistulae", "Abscess" and "Stenosis" (table 2).

Follow-up and Patient Management

Pre- and post-CTE/MRE's therapeutic plans collected from the clinical file of the patient, were analyzed for outcome evaluation. The drugs were grouped into eight categories: "Immunomodulators" (Azathioprine and Methotrexate), "Anti-TNF α agents" (Infliximab, Adalimumab and Golimumab); "Corticosteroids"; "Aminosalicylates"; "Polymeric diet"; and "Surgery". Subsequently, possible outcomes were categorized as: "No change in management", "Escalating therapy" and "De-escalating therapy"; which were subcategorized as shown in table 3.

Statistical Analysis

Descriptive analysis included demographic and clinical data were reported as a number (percent) or median (minimum, maximum, percentiles) as appropriate. The Chi-square test and Fisher's exact test were used to calculate significant associations between the age of diagnosis, disease location and disease behavior and management; the cause and the context of CTE and MRE and management; endoscopy and management; CTE/MRE findings and management; and CTE/MRE findings and specific outcomes. *P* values less than 0.05 were considered significant.

Data were analyzed by using standard statistical software, SPSS (IBM Corp. Released 2013. IBM SPSS Statistics for Windows, Version 22.0. Armonk, NY: IBM Corp.).

Ethical Considerations

This study was approved by the The Ethical Commission for Health and authorized by the board of directors of *Centro Hospitalar de São João*.

Results

A total of 349 patients were included in the study [161 were females, 188 males, with a median age of 34 years old (ranging from 11 to 76; 25 percent were 24 and 75 percent were 47) and a median age at the time of diagnosis of 24 (25 percent were 24 and 75 percent were 47)] of which 85% underwent CTE and the remaining 15% MRE. Overall, 78% of CTE or MRE were performed with the objective of monitoring the disease activity, and the remaining 22% due to the manifestation of acute disease. In 83% of cases, CTE or MRE were performed as part of a consultation, while 17% were performed during hospitalization.

The majority of cases in the study (64%) received a diagnosis between the ages of 17 and 40, presenting a non-stricturing and non-penetrating behavior (30%) and a predominantly ileal (45%) and ileocolonic (44%) disease. Baseline characteristics of the patients included in our study are presented in table 1.

Prior Endoscopic Study

Of the 349 patients in the study, 122 (36%) had undergone at least one type of endoscopic exam. Of these 122 patients, 2 were subjected only to upper digestive endoscopy, 93 to lower digestive endoscopy, and 27 to both.

CTE and MRE Reports

Radiologists reported 57 different combinations of radiological signs with an average of 2.25 findings per CTE or MRE, ranging from 0 to 6 findings. The two most common findings included bowel wall thickening and mucosal enhancement (table 2).

Of the 349 CTE or MRE performed, 60 were considered normal, of which 70% maintained the treatment plan while 30% changed it. Of the 289 abnormal exams, 42% changed the management plan while 58% maintained the treatment plan. No statistically significant association was found between a normal or abnormal exam and outcome ($p = 0.087$).

Patient Management

After CTE or MRE, 139 patients (40%) changed the management plan, and some therapeutic decisions were included in more than one category. The three most frequent alterations included the introduction of an immunomodulator agent (10%), the introduction of an anti-TNF α agent (8%) and the introduction of a corticosteroid (6%). The results related to patients' outcome are expressed in table 4.

Association between clinical factors and endoscopic study with change in management

Age at the diagnosis, disease behavior and disease location were not statistically related to changes in management (A: $p=0.121$; B: $p=0.696$; p: $p=0.090$ L1-3: $p=0.222$; L4: $p=0.812$). On the contrary, reasons for CTE and MRE and context were significantly related to the outcome (table 5).

Regarding the endoscopy performed at the time of CTE and MRE, since no statistically significant association with the outcome ($p = 0.089$) was found, its correction was not necessary for the evaluation of the role of cross-sectional imaging in the patients' outcome.

Association between specific CTE or MRE findings and change in management

Tables 6 and 7 show the results regarding the association between specific radiological signs and possible outcomes. When comparing the specific CTE and MRE features with the outcomes (change in management plan and no change in management plan), of the 70 patients with normal CTE or MRE, 18 (30%) had a change in outcome; of which 20% the outcome consisted of escalating therapy and 6% in de-escalating therapy. In our study, patients with wall thickening (74%), mucosal enhancement (65%), mural stratification (29%) and stenosis (28%) detected by the CTE or MRE were more likely to have a change in management plans. However, the only statistically significant association found was the presence of mural stratification ($p = 0.019$). Escalating therapy was more commonly associated with bowel wall thickening (74%), mucosal enhancement (69%), mural stratification (29%) and stenosis (32%), whereas bowel wall thickening (74%), mucosal enhancement (56%), mural stratification (28%) and comb sign (28%) were more often related to de-escalating therapy.

Among the changes in management plan, the presence of 2 or 3 signs is the most common combination (24% and 22%, respectively). However, 5 or 6 signs are more frequently associated with changes in the therapeutic scheme. Table 8 shows the relationship between the number of radiological signs and the outcome.

Association between specific CTE or MRE findings and specific outcomes

For patients whose outcome consisted of the introduction of an immunomodulator (table 9), the most common CTE and MRE findings were: bowel wall thickening (82%), mural hyperenhancement (79%), mural stratification (38%), presence of fistulae (27%) and presence of stenosis (27%). Mural hyperenhancement and

mural stratification on CTE and MRE were strongly related to this specific outcome (p values of 0.016 and 0.019, respectively).

The presence of ulcers, abscesses, and stenosis most commonly led to the introduction of an anti-TNF α agent (table 10). The CTE and MRE findings reported in the only case where the outcome was the increase of the dose of the immunomodulator (table 11) were mural hyperenhancement and the presence of ulcerative disease. All radiological signs of active disease showed a higher frequency for increasing the dose of anti-TNF α agent (table 12) compared to not changing the management plan, except in the presence of an abscess. In the group of patients who changed to a different immunomodulator (table 13), the CTE and MRE finding that was most frequently related with this outcome was the presence of stricturing disease. It is noteworthy that none of these associations were found to be statistically significant.

In the only case whose outcome consisted of shortening the interval between the administration of an anti-TNF α agent (table 14), only the ulcerative and structuring disease were found. However, only the association of ulcerative disease with this outcome was found to be statistically relevant ($p= 0.023$).

Discussion

CTE and MRE have an established role in evaluating patients with CD, by providing additional information to clinical and endoscopic assessment, which is crucial for proper patient management.¹⁷

Ileocolonoscopy is the gold-standard for assessment of ileocolonic CD, although the examination has certain limitations¹², such as the inability to determine disease severity in the small bowel and to evaluate extraluminal complications. CTE and MRE should represent an important step in the diagnostic, therapeutic and prognostic management of the disease in order to their ability to evaluate not only the intestinal mucosa, but also the entire bowel wall, and to detect extra luminal complications.^{18,19} CTE and MRE have a high and comparable sensitivity (85.8% vs. 87.9%) and specificity (83.6% vs. 81.2%) for active small bowel inflammation and can objectively assess response to medical therapy.^{11,13} Our study aimed to determine how specific CTE and MRE findings influenced gastroenterologists' therapeutic decisions for

patients with CD. Thereunto, we evaluated 349 patients with a previously established diagnosis of CD. In contrast to other studies, we simultaneously evaluated patients who had undergone a CTE (85% of the sample) and a MRE (15%).

The present study demonstrated that CTE and MRE alter the management plans in 40% of patients with established CD, which is in line with a previous report by a related study from Bruining et al⁷. His team verified that CTE changes the management plans in more than 50% of patients with established or suspected CD. In another study, Bruinning et al²⁰ also demonstrated that the CTE finding of penetrating disease significantly alter management plans, including the start of antibiotic therapy, surgical referral, and the use of immunosuppressive or biologic agents. Booya et al²¹ described the detection of occult penetrating disease by CTE, leading to a change in management in 61% of patients with CD. On the other hand, Figueiredo et al²² found that MRE played a decisive role in clinical decision making, leading to a change in management plans in more than half of patients. This study also confirms a previous retrospective report by Mendonza et al²³, in which MRE findings were useful for escalating medical treatment for almost 50% of patients in whom it was performed for the purpose of monitoring medical treatment.

When compared to previously studies, the patients who underwent CTE or MRE in our study were more likely to have the age at the time of diagnosis <40 years old (84%), small bowel disease (89%) and more complex disease behavior as shown by the high proportion of patients with stricturing and penetrating disease (53%). Age at the time of diagnosis <40 years old, small bowel disease and stricturing and penetrating disease (bowel damage) are some of the prognostic factors for severe disease and development of complications.²⁴ In a previous retrospective study by Mendonza et al²³ 70% of the cases were under 40 years old at the time of diagnosis, 65% had small bowel disease and 49% had stricturing or penetrating disease. However, in both studies, 24% of the cases had undergone previous surgery. In another study, Messaris et al²⁵ studied a sample of 120 patients diagnosed or suspected of CD, of which 83% were diagnosed at the age of 40, 83% had small bowel disease, and 23% were found to have stricturing or penetrating diseases. Therefore, we can assume that our sample of patients with CD was more likely to have a more complex phenotype of the disease compared with other studies. Consequently, it is not surprising that the most common outcomes in the present

study include the addition of potent anti-inflammatory medications: immunomodulators (10%), anti-TNF α agents (8%) and corticosteroids (6%).

When comparing the reason and the context of the CTE and MRE with the outcome (table 5), we found that patients who underwent CTE or MRE for CD follow-up and within the context of a consultation were more likely to have no change in their management plan (65% and 63% vs. 35% and 37%, respectively). On the contrary, for patients who underwent CTE or MRE due to acute CD complication and for those who were hospitalized, a greater proportion presented changes in the outcome (56% and 54% vs. 44% and 46%, respectively), particularly with escalating therapy (42% of 56%). These results are most likely related to the severity and acute presentation of the disease.

In the present study, 17% of the 349 CTE and MRE performed were considered normal, and, of those, 30% had a change in outcome. The remaining 83% of the patients had an abnormal CTE or MRE, of which 42% had a change in their management plan. However, unlike the study of Rajabi et al⁵ we were not able to find a statistically significant association between these two variables. In their study, they found that having an abnormal MRE was strongly associated with a change in management ($p=0.008$).

The most frequent CTE/MRE findings were bowel wall thickening (71%) and mural hyperenhancement (60%). Bowel wall thickening (bowel wall greater than 3mm in thickness) is often asymmetric, usually more prominent along the mesenteric border. Mural hyperenhancement is the most relevant finding for active disease and corresponds to a segmental attenuation greater than the adjacent normal small bowel loops.^{26,27} This CTE/MRE findings were the most commonly signs associated with change in management (74% of the patients with bowel wall thickening and 65% with mural hyperenhancement). This data is similar to a retrospective study by Rajabi et al⁵, which demonstrated that patients who had a change in management were more likely to have findings of circumferential wall thickening on MRE, as well as reactive lymph nodes. Another abnormality associated with active disease is comb sign (engorgement of *vasa recta*), but it has limited sensitivity and specificity for determining the presence of active inflammation.²⁸ In the present study, comb sign was present in 22% of the

patients and was more commonly associated with no change in management than with a change in outcome (55% and 46%, respectively, with $p>0.05$).

In addition to the signs of active disease, grading the severity of lesions is of vital importance in clinical practice since it has therapeutic and prognostic implications. The CTE and MRE findings associated to severe disease are the presence of edema, ulcerations, structuring disease and penetrating disease.²⁸ Mural stratification (figures 1 and 2) was present in 22% of the patients in the present study. This radiologic signal refers to a laminated appearance of thickened wall, due to mural edema.²⁶ It has been suggested that mural hyperenhancement, without stratification, may be more common in the early stages of CD, whereas mural stratification may indicate more advanced disease.³ Furthermore, mural edema is predominantly identified in segments with severe inflammatory lesions.²⁸ In our study, we have found that the only CTE/MRE finding that correlates significantly with change in management was mural stratification ($p<0.05$), which may be related to more advanced and severe disease in those patients. Furthermore, mural hyperenhancement and mural stratification were significantly associated with the introduction of immunomodulator ($p=0.016$ and $p=0.019$, respectively), corresponding to 79% and 38% of CTE/MRE findings associated with this outcome. Ulcerations are more commonly found on the mesenteric border of the bowel and their appearance may vary from subtle lesions (fissuring ulcerations) to deep lesions in a thickened hyperenhanced wall.²⁸ Presence of ulcerating disease was shown to lead more commonly to an escalation in therapy (50%) than to no change in the management plan (38%) or de-escalating therapy (13%).

Fibrostenosing and penetrating disease are the main complications of CD. Abscess, fistulae and stenosis are clearly signs of bowel damage, and the proportion of patients developing these complications increases over the years after a diagnosis of CD.¹⁹ Cross-sectional imaging, specifically CTE and MRE, are essential in the evaluation of these complications, since, unlike endoscopic methods, they are able to delineate fistulous pathways, detect abscesses, and identify, locate and evaluate the extent of stricturing processes.²⁸ In our study, the presence of fistulae, abscesses and stenosis were more commonly associated with the introduction of an immunomodulator and with the introduction of an anti-TNF α agent when compared to other outcomes. However, only stenosis were marginally associated with a change in management ($p=0.067$), but none of these findings were found to be strongly correlated with any

outcome ($p>0.05$). In contrast, Rajabi et al⁵ found that the presence of an abscess or phlegmon on MRE was significantly associated ($p= 0.03$) with a change in management, but they were not able to find a significant association between the presence of fistulae and a change in management.

The increased accessibility of cross-sectional imaging methods in recent years has led to an increase in the number of CT scans prescribed. Factors like young age at the time of diagnosis, the recurrent nature of the disease and the introduction of biological therapies, are related to the increased use of CTE and, consequently, to a higher cumulative radiation exposure. Thus, since MRE is a radiation-free method, it is the preferred cross-sectional imaging technique for monitoring patients with CD.²⁹ Scoring of disease activity is important for objective assessment, and several indices have been proposed. Rimola et al²⁹ proposed and validated the most widely used MRE score - MaRIA score - that includes wall thickening, mucosal hyperenhancement and presence of edema and ulcers. This index has been successfully validated using a cohort of 49 patients and ileocolonoscopy as the reference standard.¹⁴ It was demonstrated that the MaRIA score has a high accuracy for detection of active disease, severe disease, while also providing an objective, quantitative and reproducible measure of disease activity.¹⁴ Notwithstanding, this score does not appear to guide the management of CD patients and does not take into account important parameters concerning CD severity, such as the presence of stricturing disease or penetrating disease. Therefore, the development of severity scores that take these signs into account is relevant, providing important additional information about the real activity of the disease and allowing clinicians to make more informed decisions about the best therapeutic option for each case. In our study, several CTE and MRE findings were shown to be strongly related to specific outcomes: mural hyperenhancement and mural stratification with the introduction of an immunomodulator; and ulcerative disease with shortening of the interval between the administration of an anti-TNF α agent. These results lead us to believe that specific CTE and MRE findings, and perhaps new disease scoring systems, may help to stratify treatment options. However, further studies are required to validate this.

Our study highlights the importance of CTE and MRE features on the outcome of patients with CD, providing additional data about parameters that are most likely to influence clinicians' management plans of these patients. However, several limitations

of this study are evident. Given its retrospective nature, the existence of evaluation and detection biases is a possibility. Many patients had incomplete clinical data, which may have also influenced the results. Moreover, the elaboration of CTE and MRE reports by different radiologists and their evaluation and interpretation by non-radiologists may also be a factor of subjectivity and bias in the study. To prove the value of CTE and MRE in clinical assessment of CD, prospective studies are needed to determine whether the information obtained from CTE/MRE actually changes clinical decision making and clinical outcomes, and how can these imaging techniques contribute as a guide to the best therapeutic decisions.

In conclusion, the goal of our study was to retrospectively assess the impact of CTE and MRE on the therapeutic decision making for patients with CD. We found that CTE and MRE in CD patients presented important information, regarding disease activity and behavior, which played a decisive role in clinical decision making, leading to a therapeutic change in 40% of patients. Furthermore, mural stratification, a sign of CD's severity, showed to be strongly related with the outcome. These data provide additional support favoring the routine use of these imaging techniques in the monitoring and therapeutic management for patients with CD.

Acknowledgments

A work like this is the result of the involvement and collaboration of a group of people and institutions, and I would like to express my thanks:

To Professor Fernando Magro, supervisor of this dissertation, for the support and availability demonstrated throughout this study.

To the *Centro Hospitalar de São João* and to the Faculty of Medicine of the University of Porto, which allowed the concretization of this project.

To Professor Isabel Ramos for the suggestion of the project theme and indication of some bibliographical references.

To Dr. Camila Dias for the support in the statistical analysis and results interpretation.

To Dr. Rui Cunha for the availability and selection of illustrative CTE and MRE images of Crohn's Disease.

To my family and friends who have always supported me.

References

1. Torres J, Mehandru S, Colombel J et al. Crohn's disease. *The Lancet*. 2016;1 - 15. Available at: [http://dx.doi.org/10.1016/S0140-6736\(16\)31711-1](http://dx.doi.org/10.1016/S0140-6736(16)31711-1) [Accessed February 6, 2017].
2. Cheriyan D, Slattery E, McDermott S et al. Impact of magnetic resonance enterography in the management of small bowel Crohn's Disease. *Eur J Gastroenterol Hepatol*. 2013;25:550-555.
3. Allen B, Leyendecker J. MR Enterography for Assessment and Management of Small Bowel Crohn Disease. *Radiologic Clinics of North America*. 2014;52:799-810.
4. Molodecky N, Soon I, Rabi D et al. Increasing Incidence and Prevalence of the Inflammatory Bowel Diseases With Time, Based on Systematic Review. *Gastroenterology*. 2012;142:46-53.
5. Rajabi S, Dupuis I, Benmassaoud A et al. Magnetic Resonance Enterography in the Study of Patients With Crohn's Disease: Which Findings Are More Likely to Change Patient Management?. *Canadian Association of Radiologists Journal*. 2016;67:387-394.
6. Stoppino L, Della Valle N, Rizzi S et al. Magnetic resonance enterography changes after antibody to tumor necrosis factor (anti-TNF) alpha therapy in Crohn's disease: correlation with SES-CD and clinical-biological markers. *BMC Medical Imaging*. 2016;16.
7. Bruining D, Siddiki H, Fletcher J et al. Benefit of Computed Tomography Enterography in Crohn's Disease: Effects on Patient Management and Physician Level of Confidence. *Inflammatory Bowel Diseases*. 2012;18:219-225.
8. Rimola J, Alvarez-Cofiño A, Pérez-Jeldres T et al. Comparison of three magnetic resonance enterography indices for grading activity in Crohn's disease. *Journal of Gastroenterology*. 2016.
9. Higgins P, Caoili E, Zimmermann M et al. Computed tomographic enterography adds information to clinical management in small bowel Crohn's disease. *Inflammatory Bowel Diseases*. 2007;13:262-268.

10. Gomollón F, Dignass A, Annese V et al. 3rd European Evidence-based Consensus on the Diagnosis and Management of Crohn's Disease 2016: Part 1: Diagnosis and Medical Management. *Journal of Crohn's and Colitis*. 2016;11:3-25.
11. Ohtsuka K, Takenaka K, Kitazume Y et al. Magnetic resonance enterography for the evaluation of the deep small intestine in Crohn's disease. *Intestinal Research*. 2016;14:120.
12. García-Bosch O, Ordás I, Aceituno M et al. Comparison of Diagnostic Accuracy and Impact of Magnetic Resonance Imaging and Colonoscopy for the Management of Crohn's Disease. *Journal of Crohn's and Colitis*. 2016;10:663-669.
13. Bruining D, Bhatnagar G, Rimola J et al. CT and MR enterography in Crohn's disease: current and future applications. *Abdominal Imaging*. 2015;40:965-974.
14. Rimola J, Ordás I, Rodríguez S et al. Magnetic resonance imaging for evaluation of Crohn's disease. *Inflammatory Bowel Diseases*. 2011;17:1759-1768.
15. Satsangi J, Silverberg M, Vermeire S et al. The Montreal classification of inflammatory bowel disease: controversies, consensus, and implications. *Gut*. 2006;55:749-753.
16. Rieder F, Latella G, Magro F et al. European Crohn's and Colitis Organisation Topical Review on Prediction, Diagnosis and Management of Fibrostenosing Crohn's Disease. *Journal of Crohn's and Colitis*. 2016;10:873-885.
17. Panés J, Bouzas R, Chaparro M et al. Systematic review: the use of ultrasonography, computed tomography and magnetic resonance imaging for the diagnosis, assessment of activity and abdominal complications of Crohn's disease. *Alimentary Pharmacology & Therapeutics*. 2011;34:125-145.
18. Minordi L, Scaldaferri F, Larosa L et al. Comparison between clinical and radiological evaluation before and after medical therapy in patients with Crohn's disease: new prospective roles of CT enterography. *La radiologia medica*. 2014;120:449-457.
19. Church P, Turner D, Feldman B et al. Systematic review with meta-analysis: magnetic resonance enterography signs for the detection of inflammation and intestinal

damage in Crohn's disease. *Alimentary Pharmacology & Therapeutics*. 2014;41:153-166.

20. Bruining D, Siddiki H, Fletcher J et al. Prevalence of penetrating disease and extraintestinal manifestations of Crohn's disease detected with CT enterography. *Inflammatory Bowel Diseases*. 2008;14:1701-1706.

21. Booya F, Akram S, Fletcher J et al. CT enterography and fistulizing Crohn's disease: clinical benefit and radiographic findings. *Abdominal Imaging*. 2008;34:467-475.

22. Figueiredo P, Cardoso C, Ramalho M et al. P123 Impact of magnetic resonance enterography on decision-making in small-bowel Crohn's disease. *Journal of Crohn's and Colitis*. 2012;6:S59.

23. Mendoza J, González-Lama Y, Taxonera C et al. Using of magnetic resonance enterography in the management of Crohn's disease of the small intestine: First year of experience. *Revista Española de Enfermedades Digestivas*. 2012;104:578-583.

24. Peyrin-Biroulet L, Panés J, Sandborn W et al. Defining Disease Severity in Inflammatory Bowel Diseases: Current and Future Directions. *Clinical Gastroenterology and Hepatology*. 2016;14:348-354.e17.

25. Messaris E. Role of Magnetic Resonance Enterography in the Management of Crohn Disease. *Archives of Surgery*. 2010;145:471.

26. Huprich J, Fletcher J. CT enterography: Principles, technique and utility in Crohn's disease. *European Journal of Radiology*. 2009;69:393-397.

27. Kim S. Computed Tomography Enterography and Magnetic Resonance Enterography in the Diagnosis of Crohn's Disease. *Intestinal Research*. 2015;13:27.

28. Rimola J, Panés J, Ordás I. Magnetic resonance enterography in Crohn's disease: optimal use in clinical practice and clinical trials. *Scandinavian Journal of Gastroenterology*. 2014;50:66-73.

29. Rimola J, Rodriguez S, Garcia-Bosch O et al. Magnetic resonance for assessment of disease activity and severity in ileocolonic Crohn's disease. *Gut*. 2009;58:1113-1120.

Tables

Table 1. Baseline characteristics of patients with Crohn's Disease who were included in the study (CTE – Computed Tomography Enterography; MRE – Magnetic Resonance Enterography)

| | n (%) |
|-----------------------------------|--------------|
| Demographics | |
| Female | 161 (46) |
| Male | 188 (54) |
| Median age (range) | 34 (11-76) |
| Age < 18 years old | 33(10) |
| Age at diagnosis | |
| A1 | 69 (20) |
| A2 | 218 (64) |
| A3 | 54 (16) |
| Behaviour | |
| B1 | 162 (48) |
| B2 | 77 (23) |
| B3 | 102 (30) |
| P | 74 (22) |
| Location | |
| L1 | 158 (45) |
| L2 | 38 (11) |
| L3 | 152 (44) |
| L4 | 63 (18) |
| CTE's and MRE's Reason | |
| Follow-up | 271 (78) |
| Acute complication | 78 (22) |
| CTE's and MRE's context | |
| Appointment | 291 (83) |
| Internment | 58 (17) |
| History of Hospitalization | 235 (67) |
| History of Surgery | 83 (24) |

Table 2. Computed Tomography Enterography and Magnetic Resonance Enterography findings

| | n (%) |
|------------------------------|--------------|
| Bowel wall thickening | 249 (71) |
| Mucosal enhancement | 210 (60) |
| Mural stratification | 78 (22) |
| Comb sign | 77 (22) |
| Ulcer | 8 (2) |
| Fistulae | 66 (19) |
| Abscess | 11 (3) |
| Stenosis | 83 (24) |
| None | 60 (17) |

Table 3. Categorization of the outcomes in study

| No change in management | Escalating therapy | De-escalating therapy |
|--------------------------------|--|---|
| | 1. Introduction of immunomodulator | 1. Suspension of immunomodulator |
| | 2. Introduction of anti-TNF agent | 2. Suspension of anti-TNF agent |
| | 3. Dose increase of immunomodulator | 3. Dose decrease of immunomodulator |
| | 4. Dose increase of anti-TNF agent | 4. Dose decrease of anti-TNF agent |
| | 5. Shortening the interval between administrations of anti-TNF agent | 5. Extending the interval between administrations of anti-TNF agent |
| | 6. Change to a different immunomodulator | 6. Suspension of corticosteroids |
| | 7. Change to a different anti-TNF agent | 7. Decreased corticosteroid dose |
| | 8. Introduction of corticosteroid | 8. Suspension of aminosalicylates |
| | 9. Dose increase of corticosteroid | 9. Suspension of polymeric diet |
| | 10. Introduction of aminosalicylates | |
| | 11. Introduction of polymeric diet | |
| | 12. Surgery | |

Table 4. Frequency of the outcomes

| Outcome | n (%) |
|--|--------------|
| No change in management | 210 (60) |
| Escalating therapy | |
| Introduction of immunomodulator | 34 (10) |
| Introduction of anti-TNF α agent | 28 (8) |
| Dose increase of immunomodulator | 1 (0) |
| Dose increase of anti-TNF α agent | 13 (4) |
| Shortening the interval between administrations of anti-TNF α agent | 1 (0) |
| Change to a different immunomodulator | 3 (1) |
| Change to a different anti-TNF α agent | 0 (0) |
| Introduction of corticosteroid | 20 (6) |
| Dose increase of corticosteroid | 1 (0) |
| Introduction of aminosalicylates | 7 (2) |
| Introduction of polymeric diet | 0 (0) |
| Surgery | 0 (0) |
| De-escalating therapy | |
| Suspension of immunomodulator | 11 (3) |
| Suspension of anti-TNF α agent | 5 (1) |
| Dose decrease of immunomodulator | 4 (1) |
| Dose decrease of anti-TNF α agent | 0 (0) |
| Extending the interval between administrations of anti-TNF α agent | 0 (0) |
| Corticosteroids suspension | 1 (0) |
| Decreased corticosteroid dose | 18 (5) |
| Suspension of aminosalicylates | 3 (1) |
| Suspension of polymeric diet | 9 (3) |
| Suspension of immunomodulator | 4 (1) |

Table 5. Reason and context in which the Computed Tomography Enterography and Magnetic Resonance Enterography were carried out and their repercussion on the clinical management for the CD patients in study (0 – No change in management plan; 1 – Escalating therapy; 2 – De-escalating therapy; R/C– Reason/Context; Out – Outcome)

| | Outcome | | | | | | | | | <i>p</i> |
|--------------------|---------|------|------|------|-------|------|------|-------|------|--------------|
| | 0 | | | 1 | | | 2 | | | |
| | n=210 | | | n=96 | | | n=43 | | | |
| | n | %R/C | %Out | n | % R/C | %Out | n | % R/C | %Out | |
| Follow-up | 175 | 65 | 83 | 63 | 23 | 66 | 32 | 12 | 74 | 0.002 |
| Acute complication | 35 | 45 | 17 | 32 | 42 | 33 | 11 | 14 | 26 | 0.004 |
| Appointment | 184 | 63 | 88 | 73 | 25 | 76 | 35 | 12 | 81 | 0.036 |
| Internement | 26 | 46 | 12 | 23 | 40 | 24 | 8 | 14 | 18 | 0.036 |

Table 6. The impact of specific Computed Tomography Enterography and Magnetic Resonance Enterography findings on patients' outcome (0 – No change in management plan; 1 – Change in management plan; RS – Radiological Signs; Out - Outcome)

| | Outcome | | | | | | <i>p</i> |
|-----------------------|---------|-----|------|-------|-----|------|--------------|
| | 0 | | | 1 | | | |
| | n=210 | | | n=139 | | | |
| | n | %RS | %Out | n | %RS | %Out | |
| None | 42 | 70 | 20 | 18 | 30 | 13 | 0,087 |
| Bowel wall thickening | 146 | 59 | 70 | 103 | 41 | 74 | 0,355 |
| Mucosal enhancement | 120 | 57 | 57 | 90 | 43 | 65 | 0,155 |
| Mural estratification | 38 | 49 | 18 | 40 | 51 | 29 | 0,019 |
| Comb sign | 42 | 55 | 20 | 35 | 46 | 25 | 0,253 |
| Ulcer | 3 | 38 | 1 | 5 | 63 | 4 | 0,274 |
| Fistulae | 35 | 53 | 17 | 31 | 47 | 22 | 0,188 |
| Abscess | 6 | 55 | 3 | 5 | 46 | 4 | 0,759 |
| Stenosis | 44 | 53 | 21 | 39 | 47 | 28 | 0,127 |

Table 7. The impact of specific Computed Tomography Enterography and Magnetic Resonance Enterography findings on patients outcome (0 – No change in management plan; 1 – Escalating therapy; 2 – De-escalating therapy; RS – Radiological Signs; Out – Outcome)

| | Outcome | | | | | | | | | <i>p</i> |
|-----------------------|---------|-----|------|------|-----|------|------|-----|------|--------------|
| | 0 | | | 1 | | | 2 | | | |
| | n=210 | | | n=96 | | | n=43 | | | |
| | n | %RS | %Out | n | %RS | %Out | n | %RS | %Out | |
| None | 42 | 70 | 20 | 12 | 20 | 13 | 6 | 10 | 14 | 0,227 |
| Bowel wall thickening | 146 | 59 | 70 | 71 | 29 | 74 | 32 | 13 | 74 | 0,650 |
| Mucosal enhancement | 120 | 57 | 57 | 66 | 31 | 69 | 24 | 11 | 56 | 0,129 |
| Mural estratification | 38 | 49 | 18 | 28 | 36 | 29 | 12 | 15 | 28 | 0,063 |
| Comb sign | 42 | 55 | 20 | 23 | 30 | 24 | 12 | 16 | 28 | 0,455 |
| Ulcer | 3 | 38 | 1 | 4 | 50 | 4 | 1 | 13 | 2 | 0,185 |
| Fistulae | 35 | 53 | 17 | 20 | 30 | 21 | 11 | 17 | 26 | 0,338 |
| Abscess | 6 | 55 | 3 | 3 | 27 | 3 | 2 | 18 | 5 | 0,699 |
| Stenosis | 44 | 53 | 21 | 31 | 37 | 32 | 8 | 10 | 19 | 0,067 |

Table 8. Number of Computed Tomography Enterography and Magnetic Resonance Enterography findings and their repercussion on the outcome of CD patients (0 – No change in management plan; 1 – Change in management plan; RS – Radiological Signs; Out - Outcome)

| CTE/MRE findings | Outcome | | | | | | <i>p</i> |
|------------------|---------|-----|------|-------|-----|------|----------|
| | 0 | | | 1 | | | |
| | n=210 | | | n=139 | | | |
| | n | %RS | %Out | n | %RS | %Out | |
| 0 | 42 | 71 | 20 | 17 | 29 | 12 | 0,115 |
| 1 | 29 | 59 | 14 | 20 | 41 | 14 | |
| 2 | 61 | 65 | 29 | 33 | 35 | 24 | |
| 3 | 44 | 59 | 21 | 31 | 41 | 22 | |
| 4 | 22 | 52 | 11 | 20 | 48 | 14 | |
| 5 | 10 | 42 | 5 | 14 | 58 | 10 | |
| 6 | 2 | 33 | 1 | 4 | 67 | 3 | |

Table 9. Impact of specific Computed Tomography Enterography and Magnetic Resonance Enterography findings on specific outcomes – Introduction of immunomodulator (0 – No change in management plan; 1 – Change in management plan; RS – Radiological Signs; Out - Outcome)

| | Introduction of immunomodulator | | | | | | p |
|-----------------------|---------------------------------|-----|------|------|-----|------|-------|
| | 0 | | | 1 | | | |
| | n=315 | | | n=34 | | | |
| | n | %RS | %Out | n | %RS | %Out | |
| None | 59 | 98 | 19 | 1 | 2 | 3 | 0.020 |
| Bowel wall thickening | 221 | 89 | 70 | 28 | 11 | 82 | 0.135 |
| Mucosal enhancement | 183 | 87 | 58 | 27 | 13 | 79 | 0.016 |
| Mural estratification | 65 | 83 | 21 | 13 | 17 | 38 | 0.019 |
| Comb sign | 71 | 92 | 23 | 6 | 8 | 18 | 0.513 |
| Ulcer | 7 | 88 | 2 | 1 | 13 | 3 | 0.563 |
| Fistulae | 57 | 86 | 18 | 9 | 14 | 27 | 0.236 |
| Abscess | 9 | 82 | 3 | 2 | 18 | 6 | 0.291 |
| Stenosis | 74 | 89 | 24 | 9 | 11 | 27 | 0.698 |

Table 10. Impact of specific Computed Tomography Enterography and Magnetic Resonance Enterography findings on specific outcomes – Introduction of anti-TNF α agent (0 – No change in management plan; 1 – Change in management plan; RS – Radiological Signs; Out - Outcome)

| | Introduction of anti-TNF α agent | | | | | | p |
|-----------------------|---|-----|------|------|-----|------|-------|
| | 0 | | | 1 | | | |
| | n=321 | | | n=28 | | | |
| | n | %RS | %Out | n | %RS | %Out | |
| None | 52 | 87 | 16 | 8 | 13 | 29 | 0.115 |
| Bowel wall thickening | 233 | 94 | 72 | 16 | 6 | 57 | 0.083 |
| Mucosal enhancement | 194 | 92 | 60 | 16 | 8 | 57 | 0.733 |
| Mural estratification | 74 | 95 | 23 | 4 | 5 | 14 | 0.286 |
| Comb sign | 74 | 96 | 23 | 3 | 4 | 11 | 0.131 |
| Ulcer | 7 | 88 | 2 | 1 | 13 | 4 | 0.491 |
| Fistulae | 61 | 92 | 19 | 5 | 8 | 18 | 0.882 |
| Abscess | 10 | 91 | 3 | 1 | 9 | 4 | 0.605 |
| Stenosis | 76 | 92 | 24 | 7 | 8 | 25 | 0.875 |

Table 11. Impact of specific Computed Tomography Enterography and Magnetic Resonance Enterography findings on specific outcomes – Dose increase of immunomodulator (0 – No change in management plan; 1 – Change in management plan; RS – Radiological Signs; Out - Outcome)

| | Dose increase of immunomodulator | | | | | | p |
|-----------------------|----------------------------------|-----|------|-----|-----|------|-------|
| | 0 | | | 1 | | | |
| | n=348 | | | n=1 | | | |
| | n | %RS | %Out | n | %RS | %Out | |
| None | 60 | 100 | 17 | 0 | 0 | 0 | 1.000 |
| Bowel wall thickening | 249 | 100 | 72 | 0 | 0 | 0 | 0.287 |
| Mucosal enhancement | 209 | 100 | 60 | 1 | 0 | 100 | 1.000 |
| Mural estratification | 78 | 100 | 22 | 0 | 0 | 0 | 1.000 |
| Comb sign | 76 | 99 | 22 | 0 | 0 | 0 | 0.221 |
| Ulcer | 8 | 100 | 2 | 1 | 2 | 100 | 1.000 |
| Fistulae | 65 | 99 | 19 | 0 | 0 | 0 | 0.189 |
| Abscess | 11 | 100 | 3 | 0 | 0 | 0 | 1.000 |
| Stenosis | 82 | 99 | 24 | 1 | 1 | 100 | 0.238 |

Table 12. Impact of specific Computed Tomography Enterography and Magnetic Resonance Enterography findings on specific outcomes – Shortening the interval between administrations of anti-TNF α agent (0 – No change in management plan; 1 – Change in management plan; RS – Radiological Signs; Out - Outcome)

| | Shortening the interval between administrations of anti-TNF α agent | | | | | | p |
|-----------------------|---|-----|------|-----|-----|------|--------------|
| | 0 | | | 1 | | | |
| | n=348 | | | n=1 | | | |
| | n | %RS | %Out | n | %RS | %Out | |
| None | 60 | 100 | 17 | 0 | 0 | 0 | 1.000 |
| Bowel wall thickening | 249 | 100 | 72 | 0 | 0 | 0 | 0.287 |
| Mucosal enhancement | 210 | 100 | 60 | 0 | 0 | 0 | 0.398 |
| Mural estratification | 78 | 100 | 22 | 0 | 0 | 0 | 1.000 |
| Comb sign | 77 | 100 | 22 | 0 | 0 | 0 | 1.000 |
| Ulcer | 7 | 84 | 2 | 1 | 13 | 100 | 0.023 |
| Fistulae | 66 | 100 | 19 | 0 | 0 | 0 | 1.000 |
| Abscess | 11 | 100 | 3 | 0 | 0 | 0 | 1.000 |
| Stenosis | 82 | 99 | 24 | 1 | 1 | 100 | 0.238 |

Table 13. Impact of specific Computed Tomography Enterography and Magnetic Resonance Enterography findings on specific outcomes – Dose increase of anti-TNF α agent (0 – No change in management plan; 1 – Change in management plan; RS – Radiological Signs; Out - Outcome)

| | Dose increase of anti-TNF agent | | | | | | p |
|-----------------------|---------------------------------|-----|------|------|-----|------|-------|
| | 0 | | | 1 | | | |
| | n=336 | | | n=13 | | | |
| | n | %RS | %Out | n | %RS | %Out | |
| None | 59 | 98 | 18 | 1 | 2 | 8 | 0.706 |
| Bowel wall thickening | 239 | 96 | 71 | 10 | 4 | 77 | 0.765 |
| Mucosal enhancement | 200 | 95 | 60 | 10 | 5 | 77 | 0.209 |
| Mural estratification | 75 | 96 | 22 | 3 | 4 | 23 | 1.000 |
| Comb sign | 72 | 94 | 21 | 5 | 7 | 39 | 0.171 |
| Ulcer | 7 | 88 | 2 | 1 | 13 | 8 | 0.264 |
| Fistulae | 62 | 94 | 19 | 4 | 6 | 31 | 0.279 |
| Abscess | 11 | 100 | 3 | 0 | 0 | 0 | 1.000 |
| Stenosis | 79 | 95 | 24 | 4 | 5 | 31 | 0.517 |

Table 14. Impact of specific Computed Tomography Enterography and Magnetic Resonance Enterography findings on specific outcomes – Change to a different immunomodulator (0 – No change in management plan; 1 – Change in management plan; RS – Radiological Signs; Out - Outcome)

| | Change to a different immunomodulator | | | | | | p |
|-----------------------|---------------------------------------|-----|------|-----|-----|------|-------|
| | 0 | | | 1 | | | |
| | n=346 | | | n=3 | | | |
| | n | %RS | %Out | n | %RS | %Out | |
| None | 60 | 100 | 17 | 0 | 0 | 0 | 1.000 |
| Bowel wall thickening | 247 | 99 | 71 | 2 | 1 | 67 | 1.000 |
| Mucosal enhancement | 209 | 100 | 60 | 1 | 0 | 33 | 0.566 |
| Mural estratification | 77 | 99 | 22 | 1 | 1 | 33 | 0.533 |
| Comb sign | 77 | 100 | 22 | 0 | 0 | 0 | 1.000 |
| Ulcer | 8 | 100 | 2 | 0 | 0 | 0 | 1.000 |
| Fistulae | 66 | 100 | 19 | 0 | 0 | 0 | 1.000 |
| Abscess | 11 | 100 | 3 | 0 | 0 | 0 | 1.000 |
| Stenosis | 82 | 99 | 24 | 2 | 1 | 33 | 0.558 |

Figure Legends

Figure 1. Stratified appearance in Crohn's Disease. Axial and coronal contrast-enhanced Computed Tomography Enterography scan of the abdomen shows concentric wall thickening of small bowel loop with a stratified appearance indicating active disease. Also note a fistula connecting the bowel loops, a common finding in penetrating subtype of Crohn's Disease.





Figure 2. Axial contrast enhanced and T2 hASTE images of Magnetic Resonance Enterography show bowel wall thickening (double arrows), luminal fluid (*) and bowel wall stratification with mucosal and serosal avidly enhancing when compared to submucosa, in a patient with Crohn's Disease.



Anexos

Inflammatory Bowel Diseases

Instructions for Authors

SCOPE

The focus of the journal *Inflammatory Bowel Diseases*® is to bring the most current information in the clinical and basic sciences to all physicians caring for patients with inflammatory bowel diseases, and to investigators performing research in inflammatory bowel diseases and related fields. The journal publishes peer-reviewed manuscripts and review articles in basic and clinical sciences, updates on clinical trials, reviews of the current literature, editorials, and other features.

ETHICAL/LEGAL CONSIDERATIONS

A submitted manuscript must be an original contribution not previously published (except as an abstract or preliminary report), must not be under consideration for publication elsewhere, and, if accepted, must not be published elsewhere in similar form, in any language, without the consent of the publisher. Each person listed as an author is expected to have participated in the study to a significant extent. Although the editors and referees make every effort to ensure the validity of published manuscripts, the final responsibility rests with the authors, not with the journal, its editors, or the publisher. All manuscripts must be submitted through the journal's online submission system at <http://www.editorialmanager.com/ibd/>.

Inflammatory Bowel Diseases utilizes the plagiarism software, CrossCheck. Manuscripts submitted to this journal may be screened for plagiarism against previously published works. Authors who have been found to be guilty of plagiarism after the appropriate institutional investigation will be banned from submitting to the journal for a defined period of time.

Patient Anonymity and Informed Consent - It is the author's responsibility to ensure that a patient's anonymity be carefully protected and to verify that any experimental investigation with human subjects reported in the manuscript was performed with informed consent and following all the guidelines for experimental investigation with human subjects required by the institution(s) with which all the authors are affiliated. Authors should mask patients' eyes and remove patients' names from figures unless they obtain written consent from the patients and submit written consent with the manuscript.

Human Studies - Human experimentation must conform to ethical standards, and be approved by the appropriate Institutional Review Board (IRB). A statement concerning IRB approval and consent procedures must appear at the beginning of the Methods section. Any systematic data gathering effort in patients or volunteers must be approved by an IRB or adhere to appropriate local/national regulations. The Editors of *Inflammatory Bowel Diseases* are concerned about appropriate IRB review and informed consent. Authors may be questioned about the details of consent forms or the consent process. On occasion, the Editor-in-Chief may request a copy of the approved IRB application from the author. Lack of appropriate consent or documentation may be grounds for rejection. Local IRB approval does not guarantee acceptability; the final decision will be made by the Editor-in-Chief.

Animal Studies - Experimental work on animals must conform to the guidelines laid out in the Guide for the Care and Use of Laboratory Animals, which is available from the National Academy of Science; a text-only version is available at <http://www.nap.edu/readingroom/books/labrats/>. Adherence to all relevant regulations and/or approval of the appropriate institutional Animal Care Committee or governmental licensure of the investigator and/or laboratory must be obtained. A statement concerning such approval must be included at the beginning of the Methods section. The Editors of *Inflammatory Bowel Diseases* are concerned about appropriate animal care. On occasion, the Editor-in-Chief may request a copy of the approved Animal Care Committee application from the author. Local committee approval does not guarantee acceptability; the final decision will be made by the Editor-in-Chief.

Registration of Clinical Trials - All clinical trials that involve investigational drugs supported by a pharmaceutical firm or investigational devices supported by a device manufacturer must be registered at the time that a manuscript is submitted to *Inflammatory Bowel Diseases* for publication. The registry and registration number must be stated in the first paragraph of the Methods section of the manuscript.

Compliance with NIH and Other Research Funding Agency Accessibility Requirements - A number of research funding agencies now require or request authors to submit the post-print (the article after peer review and acceptance but not the final published article) to a repository that is accessible online by all without charge. As a service to our authors, LWW will identify to the National Library of Medicine (NLM) articles that require deposit and will transmit the post-print of an article based on research funded in whole or in part by the National Institutes of Health, Wellcome Trust, Howard Hughes Medical Institute, or other funding agencies to PubMed Central. The revised Copyright Transfer Agreement & Disclosure form provides the mechanism.

Copyright - All authors must complete and submit the journal's copyright transfer agreement, which includes a section on the disclosure of potential conflicts of interest based on the recommendations of the International Committee of Medical Journal Editors, "Uniform Requirements for Manuscripts Submitted to Biomedical Journals" (www.icmje.org/update.html).

A copy of the form is made available to the submitting author within the Editorial Manager submission process. Co-authors will automatically receive an Email with instructions on completing the form upon submission.

Conflicts of Interest - Authors must state all possible conflicts of interest in the manuscript, including financial, consultant, institutional and other relationships that might lead to bias or a conflict of interest. If there is no conflict of interest, this should also be explicitly stated as none declared. All sources of funding should be acknowledged in the manuscript. All relevant conflicts of interest and sources of funding should be included on the title page of the manuscript with the heading "Conflicts of Interest and Source of Funding".

For example: Conflicts of Interest and Source of Funding: A has received honoraria from Company Z. B is currently receiving a grant (#12345) from Organization Y, and is on the speaker's bureau for Organization X — the CME organizers for Company A. For the remaining authors none were declared.

In addition, each author must complete and submit the journal's copyright transfer agreement, which includes a section on the disclosure of potential conflicts of interest based on the recommendations of the International Committee of Medical Journal Editors, "Uniform Requirements for Manuscripts Submitted to Biomedical Journals" (www.icmje.org/update.html).

A copy of the form is made available to the submitting author within the Editorial Manager submission process. Co-authors will automatically receive an Email with instructions on completing the form upon submission.

Permissions - Authors must submit written permission from the copyright owner (usually the publisher) to use direct quotations, tables, or illustrations that have appeared in copyrighted form elsewhere, along with complete details about the source. Any permissions fees that might be required by the copyright owner are the responsibility of the authors requesting use of the borrowed material, and not the responsibility of the publisher.

PREPARATION OF MANUSCRIPT

Manuscripts that do not adhere to the following instructions will be returned to the corresponding author for technical revision before undergoing peer review. See specific guidelines for Letters to the Editor below. For questions on preparing manuscripts for submission, please contact Meghann Knowles at meghann@jjeditorial.com.

Title Page Include on the title page (a) complete manuscript title; (b) authors' full names, highest academic degrees, and affiliations; (c) name and address for correspondence, including fax number, telephone number, and e-mail address; and (d) sources of support that require acknowledgment.

The title page must also include disclosure of funding received for this work from any of the following organizations: National Institutes of Health (NIH); Wellcome Trust; Howard Hughes Medical Institute (HHMI); and other(s).

Structured Abstract and Key Words Limit the abstract to 250 words. It must be factual and comprehensive. Do not cite references in the abstract. Limit the use of abbreviations and acronyms, and avoid general statements (e.g., "the significance of the results is discussed"). It should be sectioned into Background, Methods, Results, and Conclusions. List three to five key words.

Text For full-length research articles, organize the manuscript in the following sequence: Abstract and Key Words, Introduction, Materials and Methods, Ethical Considerations, Results, Discussion, Acknowledgment, References, Tables, and Figure Legends. Original Research Articles: Authors are encouraged to submit articles in basic or clinical science. Review Articles of exceptional merit will also be accepted. Contributors should communicate with the Editors before submitting a review. Abbreviations must be defined at first mention in text and in each table and figure. If a brand name is cited, manufacturer and address (city and state/country) should be supplied. Acknowledge all forms of support, including pharmaceutical industry support, in an Acknowledgment paragraph.

Abbreviations For a list of standard abbreviations, consult the Council of Biology Editors Style Guide (available from the Council of Science Editors, 9650 Rockville Pike, Bethesda, MD

20814) or other standard sources. Write out the full term for each abbreviation at its first use unless it is a standard unit of measure.

References The authors are responsible for the accuracy of the references. Key the references (double-spaced) at the end of the manuscript. Cite the references in text in the order of appearance, including those references cited in tables and figure legends at the chronological citation of the tables and figures in text. Cite unpublished data, such as papers submitted but not yet accepted for publication, personal communications, in parentheses in the text. If there are more than three authors, only name the first three authors and then use et al. For abbreviations of journal names, refer to List of Journals Indexed in Index Medicus. This can be accessed at <http://www.nlm.nih.gov/tsd/serials/lji.html>.

Sample references are given below:

Journal Article

1. Gudlaugsdottir S, van Dekken H, Stijnen T, et al. Prolonged use of proton pump inhibitors, CagA status, and the outcome of *Helicobacter pylori* gastritis. *J Clin Gastroenterol*. 2002;34:536-540.

Book Chapter

2. Tobin RW, Kimmey MB. Painful diseases of the gastrointestinal tract. In: Loeser JD, ed. *Bonica's Management of Pain*. 3rd ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2001:1269-1292.

Entire Book

3. Rohen JW, Yokochi C, Lütjen-Drecoll E. *Color Atlas of Anatomy: A Photographic Study of the Human Body*. 5th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2002.

Software

4. Epi Info [computer program]. Version 6. Atlanta: Centers for Disease Control and Prevention; 1994.

Online Journals

5. Friedman SA. Preeclampsia: a review of the role of prostaglandins. *ObstetGynecol* [serial online]. January 1988;71:22-37. Available from: BRS Information Technologies, McLean, VA. Accessed December 15, 1990.

Database

6. CANCERNET-PDQ [database online]. Bethesda, MD: National Cancer Institute; 2014. Updated March 29, 2014.

World Wide Web

7. Gostin LO. Drug use and HIV/AIDS [JAMA HIV/AIDS Web site]. June 1, 2015. Available at: <http://www.ama-assn.org/special/hiv/ethics>. Accessed July 26, 2015.

Figures Art should be created/scanned and saved and submitted as either a TIFF (tagged image file format), or an EPS (encapsulated postscript) file. Line art must have a resolution of at least 1200 dpi (dots per inch), and electronic photographs, radiographs, CT scans, and other scanned images must have a resolution of at least 300 dpi. If fonts are used in the artwork, they must be converted to paths or outlines or they must be embedded in the files. Color images must be created/scanned and saved and submitted as CMYK files. Please note that artwork generated from office suite programs such as Corel Draw and MS Word and artwork downloaded from the Internet (JPEG or GIF files) cannot be used. Cite figures consecutively in the manuscript, and number them in the order in which they are discussed.

Figure Legends Legends must be submitted for all figures. They should be brief and specific, and they should appear after the tables. Use scale markers in the image for electron micrographs, and indicate the type of stain used.

Color Figures At no cost to the author, the journal accepts for publication color figures that will enhance the article. Figures are also published online in color at no cost.

Steps for Submitting Artwork

1. Please reference "5 Steps to Creating Digital Artwork (pdf)" available at <http://edmgr.ovid.com/lww-final/accounts/5StepsforArt.pdf>, specifically steps 2 and 3, when creating figures for submission.
2. Create, Scan and Save according to the "5 Steps to Creating Digital Artwork (pdf)".
3. Compare your final figure to the Target Digital-Imaging Results listed below.
4. Upload each figure to the journal's online submission system in conjunction with your manuscript text and tables.

Digital Artwork Guideline Checklist

Here are the basics to have in place before submitting your digital art to *Inflammatory Bowel Diseases*. Artwork saved as TIFF and EPS files. Do not save TIFFs as compressed files. PowerPoint files are also acceptable.

- Artwork created as the actual size (or slightly larger) it will appear in the journal. (To get an idea of the size images should be when they print, study a copy of the journal to which you wish to submit. Measure the artwork typically shown and scale your image to match.)
- Crop out any white or black space surrounding the image.
- Text and fonts in any figure are one of the acceptable fonts: Helvetica, Times Roman, Symbol, Mathematical PI, and European PI.
- Color images are created/scanned and saved and submitted as CMYK only. Do not submit any figures in RGB mode because RGB is the color mode used for screens/monitors and CMYK is the color mode used for print.
- Line art saved at a resolution of at least 1200 dpi.
- Images saved at a resolution of at least 300 dpi.

- Each figure saved as a separate file and saved separately from the accompanying text file.
- For multi-panel or composite figures only: Any figure with multiple parts should be sent as one file with each part labeled the way it is to appear in print.

Remember:

- Artwork generated from office suite programs such as CorelDRAW, MS Word, Excel, and artwork downloaded from the Internet (JPEG or GIF files) cannot be used because the quality is poor when printed.
- Cite figures consecutively in your manuscript.
- Number figures in the figure legend in the order in which they are discussed.
- Upload figures consecutively to the journal's online submission system and number figures consecutively in the Description box during upload.

Tables Create tables using the table creating and editing feature of your word processing software (e.g., Word). Do not use Excel or comparable spreadsheet programs. Do not submit tables as image files or images placed in Word documents, tables must be provided as editable text (Word files are preferred). Submit all tables as separate files. Cite tables consecutively in the text, and number them in that order. Key each on a separate sheet, and include the table title, appropriate column heads, and explanatory legends (including definitions of any abbreviations used). Do not imbed tables within the body of the manuscript. They should be self-explanatory and should supplement, rather than duplicate, the material in the text.

Supplemental Digital Content (SDC) Authors may submit supplementary materials via the journal's online submission system that enhance their article's text to be considered for online posting. SDC may include standard media such as text documents, graphs, audio, video, etc. On the Attach Files page of the submission process, please select Supplemental Audio, Video, or Data for your uploaded file as the Submission Item. If an article with SDC is accepted, our production staff will create a URL with the SDC file. The URL will be placed in the call-out within the article. SDC files are not copy-edited by LWW staff, they will be presented digitally as submitted. Please supply the SDC files as you would like them to appear in final publication (include legends in the same file as the images; make text double spaced or single spaced per your preference). For a list of all available file types and detailed instructions, please visit <http://links.lww.com/A142>.

SDC Call-outs Supplemental Digital Content must be cited consecutively in the text of the submitted manuscript. Citations should include the type of material submitted (Audio, Figure, Table, etc.), be clearly labeled as "Supplemental Digital Content," include the sequential list number, and provide a description of the supplemental content. All descriptive text should be included in the call-out as it will not appear elsewhere in the article. Example: We performed many tests on the degrees of flexibility in the elbow (see Video, Supplemental Digital Content 1, which demonstrates elbow flexibility) and found our results inconclusive.

List of Supplemental Digital Content A listing of Supplemental Digital Content must be submitted at the end of the manuscript file. Include the SDC number and file type of the Supplemental Digital Content. This text will be removed by our production staff and not be published.

Example:

Supplemental Digital Content 1.wmv

SDC Files Requirements All acceptable file types are permissible up to 10 MBs. For audio or video files greater than 10 MBs, authors should first query the journal office for approval. For a list of all available file types and detailed instructions, please visit <http://links.lww.com/A142>.

Style Follow American Medical Association Manual of Style (10th edition). Stedman's Medical Dictionary (27th edition) and Merriam Webster's Collegiate Dictionary (10th edition) should be used as standard references. Refer to drugs and therapeutic agents by their accepted generic or chemical names, and do not abbreviate them. Use code numbers only when a generic name is not yet available. In that case, supply the chemical name and a figure giving the chemical structure of the drug. Capitalize the trade names of drugs and place them in parentheses after the generic names. To comply with trademark law, include the name and location (city and state in USA; city and country outside USA) of the manufacturer of any drugs, supplies, or equipment mentioned in the manuscript. Use the metric system to express units of measure and degrees Celsius to express temperatures, and use SI units rather than conventional units.

Basic and Clinical IBD Review Articles Guidelines Basic and clinical IBD Review articles should present recent advances in a relatively narrow topic that have been made in cutting edge research.

The goal of the basic and clinical IBD review articles should be to present a complete summary of important research areas that are now improving our understanding of Crohn's disease and ulcerative colitis.

The body of basic and clinical IBD Review articles should be no longer than 20 double spaced pages (not including references, figures, and tables). There should be no more than 6 tables and figures (combined). Supplemental figures and tables will be allowed online. If page limits need to be increased, the authors may request permission from the Editors to increase the length of the Review article. The number of references should be limited to 100. The Review article should be focused on a single specific topic. All review articles will be peer-reviewed.

Future Directions and Methods for IBD Research Guidelines Original research articles and review articles on future directions and methods for IBD research should discuss important basic and clinical areas in which investigators should focus their efforts to provide a deeper understanding of IBD research areas in which rapid advances and novel concepts can be made. In addition, the Future Directions and Methods for IBD original research articles and review articles should include a discussion of areas in which improved methodologic tools have been identified and exactly how those methods should be performed. Basic science directions and methods should be presented that will expand our knowledge of areas that will allow novel insights to be made regarding the genetic, immunologic, microbial and environmental interactions that are the basis of the pathogenesis of Crohn's disease and ulcerative colitis. Clinical directions and methods should be presented that will allow investigators to make advances using cohort studies, multicenter registries, risk stratifications, and treatment outcomes. The goal of the Future Directions and Methods for clinical IBD original research articles and review articles should be to better understand the challenging biological variables in IBD patients and to provide optimal evidence of novel

therapeutics that will more effectively treat, cure, and prevent Crohn's disease and ulcerative colitis.

Original research articles, for the Future Directions and Methods for IBD research section, should be written as described in the IBD Journal Instructions for original research articles. Review articles, for the Future Directions and Methods for IBD research section, should be no longer than 35 double spaced pages (including references and figure legends). There should be no more than 6 tables and figures (combined). Supplemental figures and tables will be allowed online. If page limits need to be increased, the authors may request permission from the Editors to increase the length of their future directions and methods for IBD research review article. The number of references should be limited to 100. The future directions and methods for IBD research review article should be focused on a single specific topic.

Letters to the Editor Guidelines Letters in response to articles published in the journal are welcome and should be submitted via IBD's online submission system. All Letters should start with the phrase "To the Editors," and be written as a letter. Letters must be submitted the end of the following calendar month (e.g. by the end of July, for letters referring to articles in the June print issue) and be a maximum of 400 words, with no more than one figure or table and no more than 5 references. All LTEs will be published online-only.

Research letters and case-based discussions will no longer be accepted for submission effective November 1, 2012.

REVISIONS

When submitting a revision, please submit both a clean copy and marked copy of the manuscript. The marked copy should highlight all of the changes made by authors after the original review. Authors can use the track changes feature of the Microsoft Word program to create a marked copy. Authors also should submit all tables and figures in separate files for production purposes.

AFTER ACCEPTANCE

Open access LWW's hybrid open access option is offered to authors whose articles have been accepted for publication. With this choice, articles are made freely available online immediately upon publication. Authors may take advantage of the open access option at the point of acceptance to ensure that this choice has no influence on the peer review and acceptance process. These articles are subject to the journal's standard peer-review process and will be accepted or rejected based on their own merit.

Authors of accepted peer-reviewed articles have the choice to pay a fee to allow perpetual unrestricted online access to their published article to readers globally, immediately upon publication. The article processing charge for *Inflammatory Bowel Diseases* is \$3,000. The article processing charge for authors funded by the Research Councils UK (RCUK) is \$3,800. The publication fee is charged on acceptance of the article and should be paid within 30 days by credit card by the author, funding agency or institution. Payment must be received in full for the article to be published open access.

- **Authors retain copyright** Authors retain their copyright for all articles they opt to publish open access. Authors grant LWW a license to publish the article and identify itself as the original publisher.

- **Creative Commons license** Articles opting for open access will be freely available to read, download and share from the time of publication. Articles are published under the terms of the Creative Commons License Attribution-NonCommercial No Derivative 3.0 which allows readers to disseminate and reuse the article, as well as share and reuse of the scientific material. It does not permit commercial exploitation or the creation of derivative works without specific permission. To view a copy of this license visit: <http://creativecommons.org/licenses/by-nc-nd/3.0>.
- **Compliance with NIH, RCUK, Wellcome Trust and other research funding agency accessibility requirements** A number of research funding agencies now require or request authors to submit the post-print (the article after peer review and acceptance but not the final published article) to a repository that is accessible online by all without charge. As a service to our authors, LWW identifies to the National Library of Medicine (NLM) articles that require deposit and transmits the post-print of an article based on research funded in whole or in part by the National Institutes of Health, Howard Hughes Medical Institute, or other funding agencies to PubMed Central. The revised Copyright Transfer Agreement provides the mechanism. LWW ensures that authors can fully comply with the public access requirements of major funding bodies worldwide. Additionally, all authors who choose the open access option will have their final published article deposited into PubMed Central. RCUK and Wellcome funded authors can choose to publish their paper as open access with the payment of an article process charge (gold route), or opt for their accepted manuscript to be deposited (green route) into PMC with an embargo. With both the gold and green open access options, the author will continue to sign the Copyright Transfer Agreement (CTA) as it provides the mechanism for LWW to ensure that the author is fully compliant with the requirements. After signature of the CTA, the author will then sign a License to Publish where they will then own the copyright. Those authors who wish to publish their article via the gold route will be able to publish under the terms of the Attribution 3.0 (CCBY) License. To view of a copy of this license visit: <http://creativecommons.org/licenses/by/2.0/>. Those authors who wish to publish their article via the green route will be able to publish under the rights of the Attribution Non-commercial 3.0 (CCBY NC) license (<http://creativecommons.org/licenses/by-nc/2.0/>). It is the responsibility of the author to inform the Editorial Office and/or LWW that they have RCUK funding. LWW will not be held responsible for retroactive deposits to PMC if the author has not completed the proper forms.

FAQ for open access <http://links.lww.com/LWW-ES/A48>

Page Proofs and Corrections - Corresponding authors will receive an e-mail containing a link to the electronic page proofs to check the copyedited and typeset article before publication. The pages proofs are provided as portable document format (PDF) files which require Adobe Reader to be viewed and edited. Complete instructions will be provided with the e-mail for downloading the files and for returning the corrected pages electronically to the publisher. It is the author's responsibility to ensure that there are no errors in the proofs. Changes that have been made to conform to journal style should be allowed to stand if they do not alter the authors' meaning. Authors may be charged for alterations to the proofs beyond those required to correct errors or to answer queries. Proofs must be checked

carefully and corrections returned within 24 to 48 hours of receipt, as requested in the communication accompanying the page proofs.

Reprints - The corresponding author will receive the reprint order form via e-mail while the article is in production. Reprint requests can be submitted any time before or after publication of the article. For orders submitted before the article is published, the reprints are normally shipped 6 to 8 weeks after publication of the issue in which the item appears. For any questions regarding reprints or publication fees, contact the Reprint Department by mail at Lippincott Williams & Wilkins, 351 West Camden Street, Baltimore, MD 21201; by phone at 1-800-341-2258; by fax at 410-528-4434; or by e-mail at: reprints@wolterskluwer.com.

Permissions - For permission and/or rights to use content for which the copyright holder is LWW or the society, please go to the journal's website and after clicking on the relevant article, click on the "Request Permissions" link under the "Article Tools" box that appears on the right side of the page. Alternatively, send an e-mail to customercare@copyright.com.

For Translation Rights & Licensing queries, contact Silvia Serra, Translations Rights, Licensing & Permissions Manager, Wolters Kluwer Health (Medical Research) Ltd, 250 Waterloo Road, London SE1 8RD, UK. Phone: +44 (0) 207 981 0600. E-mail: silvia.serra@wolterskluwer.com

For Special Projects and Reprints (U.S./Canada), contact Alan Moore, Director of Sales, Lippincott Williams & Wilkins, Two Commerce Square, 2001 Market Street, Philadelphia, PA 19103. Phone: 215-521-8638. E-mail: alan.moore@wolterskluwer.com

For Special Projects and Reprints (non-U.S./Canada), contact Silvia Serra, Translations Rights, Licensing & Permissions Manager, Wolters Kluwer Health (Medical Research) Ltd, 250 Waterloo Road, London SE1 8RD, UK. Phone: +44 (0) 207 981 0600. E-mail: silvia.serra@wolterskluwer.com